arXiv:1209.5784v1 [math.DS] 25 Sep 2012

Pesin's Entropy Formula for C^1 Diffeomorphisms with Dominated Splitting

Eleonora Catsigeras, Marcelo Cerminara and Heber Enrich *

September 27, 2012

Abstract

For any C^1 diffeomorphism with dominated splitting we consider a nonempty set of invariant measures which describes the asymptotic statistics of Lebesgue-almost all the orbits. They are the limits of convergent subsequences of averages of the Dirac's deltas supported on those orbits. We prove that the metric entropy of each of these measures is lower bounded by the sum of the Lyapunov exponents on the dominating subbundle. As a consequence, if those exponents are non negative, and if the exponents on the dominated subbundle are non positive, those measures satisfy the Pesin's Formula of the Entropy.

1 Introduction

As pointed out by [P84] and other authors, there is a gap between the $C^{1+\alpha}$ and the C^1 Pesin's theory. To find new results that hold for $C^1 \setminus C^{1+\alpha}$ maps, relatively recent research started assuming some uniformly dominated condition (see [**?**, **?**]).

Let us consider $f \in \mathrm{Diff}^1(M)$, where M is a compact and connected riemannian manifold of finite dimension. We denote by $\mathcal{P}$ the set of all the Borel probability measures endowed with the weak* topology, and by $\mathcal{P}_f \subset \mathcal{P}$ the set of f-invariant probabilities. We denote by m the Lebesgue measure (after rescaling so that $m \in \mathcal{P}$). For all $\mu \in \mathcal{P}_f$ for μ-a.e. $x \in M$ the Lyapunov exponents of the orbit of x are denoted by

$$\chi_1(x) \geq \chi_2(x) \geq \ldots \geq \chi_{\dim M}(x).$$

Let

$$\chi_i^+(x) := \max\{\chi_i(x), 0\}.$$

Theorem (Ruelle's Inequality) [R78]
For all $f \in \mathrm{Diff}^1(M)$ and for all $\mu \in \mathcal{P}_f$

$$h_\mu \leq \int \sum_{i=1}^{\dim M} \chi_i^+ \, d\mu,$$

*All authors: Instituto de Matemática y Estadística Rafael Laguardia (IMERL), Facultad de Ingeniería, Universidad de la República, Uruguay. E-mails: eleonora@fing.edu.uy, cerminar@fing.edu.uy, enrich@fing.edu.uy. Address: Julio Herrera y Reissig 565. Montevideo. Uruguay.

where h_μ denotes the metric theoretical entropy of μ.
Definition Let $f \in \text{Diff}^1(M)$ and $\mu \in \mathcal{P}_f$. We say that μ satisfies the *Pesin's Entropy Formula*, and write $\mu \in PF$, if

$$h_\mu = \int \sum_{i=1}^{\dim M} \chi_i^+ \, d\mu.$$

We denote by m^u the Lebesgue measure along the unstable manifolds of the regular points for which positive Lyapunov exponents exist. Also, for any regular point x for which all the Lyapunov exponents are non positive, we call unstable manifold (of dimension zero) of x to $\{x\}$, and in this case we denote to $m^u = \delta_x$.

For any invariant measure μ we denote by μ^u the conditional measures of μ along the unstable manifolds, after applying the local Rohlin decomposition [R62].

The following are well known results of the Pesin's theory under the hypothesis $f \in \text{Diff}^{\,2}(M)$

Pesin Theorem [**?**, **?**] *Let $\mu \in \mathcal{P}_f$ be hyperbolic (namely $\chi_i(x) \neq 0 \ \forall \ i \ \mu-a.e.$ $x \in M$). If $\mu \ll m$ then $\mu^u \ll m^u$ and $\mu \in PF$.*

Ledrappier-Young Theorem [LY85]: *If $\mu \in PF$ then $\mu^u \ll m^u$.*

Pugh-Shub Theorem [PS89]: *Let $\mu \in \mathcal{P}_f$ be ergodic, hyperbolic and satisfying $\mu^u \ll m^u$. Then μ is SRB* (namely, μ is physical, according to Definition 1.3).

This last result states that μ describes the asymptotic statistics of a Lebesgue-positive set of orbits (see Definition 1.4).

Many results of the Pesin's Theory, among them Ledrappier-Young Theorem, fail if $f \in C^1 \setminus C^{1+\alpha}$ (see for example [CQ01]). The general purpose of this paper is to look for adequate reformulations of some of the above results, dealing with measures $\mu \in PF$ to hold for all $f \in \text{Diff}^1(M)$ (note that C^1 generically $f \notin \text{Diff}^{1+\alpha}(M)$). To do that we recall some definitions and previous results taken from [CE11].

Definition 1.1 (Asymptotic statistics of each orbit)
Fix $x \in M$. The *sequence of empirical probabilities* of x is $\{\sigma_{n,x}\}_{n\geq 1} \subset \mathcal{P}$, where

$$\sigma_{n,x} := \frac{1}{n} \sum_{j=0}^{n-1} \delta_{f^j(x)}.$$

The *$p\omega$-limit* of x is

$$p\omega(x) := \{\mu \in \mathcal{P} : \ \exists \ n_j \to +\infty \text{ such that } \lim_{j\to\infty} \sigma_{n_j,x} = \mu\} \subset \mathcal{P}_f$$

We say that $p\omega(x)$ describes *the asymptotic statistics* of the orbit of x.

Definition 1.2 (Basins of statistical strong and weak attraction)
For any given $\mu \in \mathcal{P}$ the *basin of (strong) statistical attraction of μ* is

$$B(\mu) := \{x \in M : \ p\omega(x) = \{\mu\}\}.$$

Fix any metric dist^* in $\mathcal{P}$ endowing the weak* topology. Take $\mu \in \mathcal{P}$.

The *basin of ε-weak statistical attraction of μ* is

$$B_\varepsilon(\mu) := \{x \in M : \ \text{dist}^*(p\omega(x), \mu) < \varepsilon\}\,.$$

Definition 1.3 (SRB and SRB-like measures)

A probability measure μ is *SRB or physical* if $m(B(\mu)) > 0$. A probability measure μ is *SRB-like or pseudo-physical* if $m(B_\varepsilon(\mu)) > 0 \ \ \forall\ \varepsilon > 0$. We write $\mu \in$ SRB ($\mu \in$ SRB-like) if μ is SRB (resp. SRB-like).

It is standard to check that the set of SRB-like measures is independent of the metric dist^* that was chosen in $\mathcal{P}$ and that it is contained in $\mathcal{P}_f$.

Definition 1.4 (Description of the asymptotic statistics of the system)

Given $f\colon M \to M$, we say that a weak*-compact set $\mathcal{K} \subset \mathcal{P}$ *describes the asymptotic statistics* of Lebesgue-almost all the orbits of f if $p\omega(x) \subset \mathcal{K}$ for Lebesgue-almost all $x \in M$.

Theorems 1.3 and 1.5 of [CE11] proved that SRB-like $\neq \emptyset$, $p\omega(x) \subset$ SRB-like for Lebesgue-almost all $x \in M$, and SRB-like is the minimal weak*-compact set $\mathcal{K} \subset \mathcal{P}$ such that $p\omega(x) \subset \mathcal{K}$ for Lebesgue-almost all $x \in M$. In this sense the set SRB-like minimally describes the asymptotic statistics of Lebesgue-almost all the orbits.

Our focus is to find relations between:

- SRB or physical measures and, more generally, SRB-like measures.
- Invariant measures μ that satisfy the Pesin's Entropy Formula ($\mu \in PF$).

Two interesting results were recently obtained for $f \in \text{Diff}^1(M) \setminus \text{Diff}^{1+\alpha}(M)$:

Qiu Theorem [Q11]: *If f is a transitive Anosov, then C^1-generically there exists a unique $\mu \in PF$. Besides, μ is SRB.*

Sun-Tian Theorem [ST12]: *If $f \in \text{Diff}^1(M)$ has an invariant measure $\mu \ll m$, and if there exists a dominated splitting $E \oplus F$ such that either $\chi_F \geq 0$ and $\chi_E < 0$ or $\chi_F > 0$ and $\chi_E \leq 0$, then $\mu \in PF$.*

To prove Sun-Tian Theorem the authors use a technique introduced by Mañé [M81] (which we call "the technique of the dispersion of C^1 Hadarmard graphs"). With this method, the Pesin's Entropy Formula can be obtained without using absolute continuity properties of invariant foliations.

In this paper, we reformulate the technique of Mañé [M81], to generalize the result of [ST12] to all $f \in \text{Diff}^1(M)$ with dominated splitting, even if no invariant measure μ satisfies $\mu \ll m$. Precisely:

Definition (Dominated splitting) Let $f\colon M \to M$ be a C^1 diffeomorphism on a compact riemannian manifold. Let $TM = E \oplus F$ be a continuous and df-invariant splitting. We call $TM = E \oplus F$ to be a *dominated splitting* if there exist $C > 0$ and $0 < \lambda < 1$ such that

$$\|df^n|_{E_x}\| \left\| df^{-n}|_{F_{f^n(x)}} \right\| \leq C\lambda^n,\ \forall x \in \mathbb{N} \text{ and } n \geq 1.$$

We prove the following result:

Theorem 1 *Let $f \in Diff^{\,1}(M)$ have a dominated splitting $TM = E \oplus F$.*

Then:

(A) *There exist measures μ characterized by the following properties:*

(A1) *The set of such measures minimally describe the asymptotic statistics of Lebesgue-almost all the orbits.*

(A2) *They satisfy the inequality:*

$$h_\mu(f) \geq \int \sum_{i=1}^{\dim F} \chi_i \, d\mu \tag{1}$$

(B) *Moreover, Property* **(A1)** *implies Inequality* **(A2)** .

(C) *If besides $\chi_{\dim F} \geq 0 \geq \chi_{\dim F+1}$, then any such measure μ satisfies the Pesin's Entropy Formula.*

The inequality **(A2)**, together with with Ruelle's Inequality, implies Assertion **(C)**. Besides, Definition 1.3 of SRB-like measures implies their existence and that the set of all of them satisfies Property **(A1)** (see Theorems 1.3 and 1.5 of [CE11]). So, to prove Theorem 1 it is only left to prove its part **(B)**, namely, that any SRB-like measure satisfies Inequality **(A2)**. In other words, Theorem 1 is reduced to the following statement, whose proof is the purpose of this paper:

Theorem 2 *Let $f \in Diff^{\,1}(M)$ have a dominated splitting $TM = E \oplus F$. Let μ be a SRB-like measure for f. Then, μ satisfies Inequality* (1).

The same arguments of the proof of Theorem 2 also hold in a more general context for $f \in \text{Diff}^{\,1}(M)$. In fact, if $\Lambda \subset M$ is an invariant and compact topological attractor, and if $\overline{V} \supset \Lambda$ is a compact neighborhood with dominated splitting $T_{\overline{V}} = E \oplus F$, then the same stament of Theorems 1 and 2 are valid for $f|_V$.

Observe that the converse statement of Theorem 2 is false: consider a C^2 non transitive hyperbolic attractor, with a finite set $SRB = \{\mu_1, \mu_2, \ldots, \mu_k\}$ $(k \geq 2)$, such that SRB attracts statistically Lebesgue-almost all the orbits. Therefore, SRB-like = SRB. So, $(\mu_1 + \mu_2)/2 \notin SRB$ (we are assuming that $\mu_1 \neq \mu_2$). After Theorem 2, $\mu_1, \mu_2 \in$ PF. It is well known that any convex combination of measures in PF is also in PF (see Theorem 5.3.1 and Lemma 5.2.2. of [K98]). So $(\mu_1 + \mu_2)/2 \in$ PF but it is not SRB-like.

The paper is organized as follows: In Section 2 we reduce the proof of Theorem 2 to Lemmas 2.1 and 2.2. In Sections 3 and 4 we prove Lemmas 2.1 and 2.2 respectively. Finally, in Section 5 we show some technical assertions that are used in the proofs of the previous sections.

2 Reduction of the proof of Theorem 2

Notation:

Let $\mathcal{B}$ be the Borel-sigma algebra on the manifold M. We denote by $\alpha = \{X_i\}_{1 \leq i \leq k}$ a finite partition of M, namely, $k \in \mathbb{N}$, $X_i \in \mathcal{B}$ $\forall$ $1 \leq i \leq k$, $X_i \bigcap X_j = \emptyset$ if $i \neq j$, and $\bigcup_{i=1}^{k} X_i = M$. We write $f^{-j}(\alpha) = \{f^{-j}(X_i)\}_{1 \leq i \leq k}$.

For any pair of finite partitions $\alpha = \{X_i\}_{1\le i\le k}$ and $\beta = \{Y_j\}_{1\le j\le h}$ we write $\alpha \bigvee \beta = \{X_i \bigcap Y_j : 1 \le i, j \le k, h, X_i \bigcap Y_j \neq \emptyset\}$. We denote: $\alpha^n = \bigvee_{j=0}^{n} f^{-j}(\alpha)$.

For the diffeomorphism $f : M \mapsto M$ with dominated splitting $E_x \oplus F_x = T_xM$, we denote:

$$\psi(x) := -\log \left| \det df(x)|_{F_x} \right| \tag{2}$$

$$\psi_n(x) := -\log \left| \det df^n(x)|_{F_x} \right| = \sum_{j=0}^{n-1} \psi \circ f^j(x) = -\log \left| \det df^{-n}(f^n(x))|_{F_{f^n(x)}} \right| \tag{3}$$

Lemma 2.1 (Upper bound of the Lebesgue measure m)

For all $\varepsilon > 0$ there exists $\delta > 0$ such that for every finite partition α with $\operatorname{diam}(\alpha) < \delta$ *there exist a sequence $\{\nu_n\}_{n\ge 0}$ of finite measures and a constant $K > 0$ such that:*

(i) $\nu_n(X) < K \ \forall\ X \in \alpha^n = \bigvee_{j=0}^{n} f^{-j}(\alpha) \ \ \forall\ n \ge 0$.

(ii) $\forall\ C \in \mathcal{B}$ *and* $\forall\ n \in \mathbb{N}$:

$$m(C) \le K e^{n\varepsilon} I(\psi_n, C, \nu_n), \quad \textit{where}$$

$$I(\psi_n, C, \nu_n) := \int_C e^{\psi_n}\, d\nu_n.$$

We prove Lemma 2.1 in Section 3. Before stating the second lemma, recall Definition 1.2 of the basin $B_\epsilon(\mu)$.

Lemma 2.2 (Lower bound for the metric entropy)

For all $\mu \in \mathcal{P}_f$ and for all $\varepsilon,\ \delta > 0$ there exist a finite partition α satisfying $\operatorname{diam}\alpha < \delta$ *and a sequence $\{C_n\}_{n\ge 0} \subset \mathcal{B}$ such that:*

(iii) $\bigcap_{N\ge 0} \bigcup_{n\ge N} C_n \supset B_{\varepsilon^*}(\mu)$ *for some $\varepsilon^* > 0$ for some* dist^* *in $\mathcal{P}$.*

(iv) $\forall$ *sequence $\{\nu_n\}_{n\ge 0}$ of finite measures such that $\exists\ K > 0$ verifying $\nu_n(X) < K\ \forall\ X \in \alpha^n\ \forall\ n \ge 0$, the following inequality holds:*

$$\limsup_{n\to+\infty} \frac{1}{n} \log I(\psi_n, C_n, \nu_n) \le \varepsilon + h_\mu(\alpha) + \int \psi\, d\mu.$$

We prove Lemma 2.2 in Section 4. Now, let us show that the two lemmas above easily imply the statement of Theorem 2:

Proof of Theorem 2 assuming Lemmas 2.1 and 2.2

Let $\mu \in \mathcal{P}_f$ such that $h_\mu(f) - \int \sum_{i=1}^{\dim F} \chi_i\, d\mu = -r < 0$. We must prove that $\mu \notin$ SRB-like.

Take $0 < \varepsilon < r/4$. Consider $\delta > 0$ as given by Lemma 2.1. Construct the partition α and the sequence $\{C_n\}_{n\ge 0} \subset \mathcal{B}$ by Lemma 2.2. Construct the sequence $\{\nu_n\}_{n\ge 0}$ of finite measures and the constant $K > 0$, by Lemma 2.1.

We now apply Lemma 2.2:

$$\limsup_{n\to\infty} \frac{1}{n} \log I(\psi_n, C_n, \nu_n) \le \varepsilon + h_\mu(\alpha) + \int \psi\, d\mu \le \frac{r}{4} - r. \tag{4}$$

Besides, by Lemma 2.1:

$$\frac{1}{n}\log m(C_n) \leq \frac{\log K}{n} + \frac{r}{4} + \frac{1}{n}\log I(\psi_n, C_n, \nu_n) \text{ (recall that } 0 < \varepsilon < r/4). \qquad (5)$$

We join the two inequalities (4) and (5) to deduce that:

$$\limsup_{n\to\infty} \frac{1}{n}\log m(C_n) \leq \frac{r}{4} + \frac{r}{4} - r = -\frac{r}{2} < 0.$$

By Borel-Cantelli Lemma, $m\left(\bigcap_{N\geq 0}\bigcup_{n\geq N} C_n\right) = 0$. Using the property **(iii)** of Lemma 2.2 we obtain:

$$m(B_{\varepsilon^*}(\mu)) \leq m\left(\bigcap_{N\geq 0}\bigcup_{n\geq N} C_n\right) = 0.$$

Then μ does not satisfy Definition 1.3, i.e. it is not SRB-like. □

3 Proof of Lemma 2.1

To prove Lemma 2.1 we will use, after some reformulations, the technique of the dispersion of Hadamard graphs, introduced by Mañé in [M81].

Notation: First take a fixed value of $\delta > 0$ small enough such that $\exp_x^{-1}$ is a diffeomorphism from $B_{3\delta}(x)$ onto its image in $T_xM \ \ \forall\, x \in M$. Fix $x \in M$. Denote

$$\mathbf{B}_\delta^H(\mathbf{0}) := \{v \in H : \|v\| \leq \delta\}, \text{ where } H \text{ stands for } E_x, F_x,$$

$$\mathbf{B}_\delta^{T_xM}(\mathbf{0}) := \mathbf{B}_\delta^{E_x}(\mathbf{0}) \oplus \mathbf{B}_\delta^{F_x}(\mathbf{0}).$$

Denote by π_{E_x} (π_{F_x}) the projection of T_xM on E_x along F_x (resp. on F_x along E_x), and $\gamma := \max_{x\in M}\{\|\pi_{E_x}\|, \|\pi_{F_x}\|\}$.

$\forall\ v \in \mathbf{B}_\delta^{T_xM}(\mathbf{0})$ we write $v_1 := \pi_{E_x} v, \quad v_2 := \pi_{F_x} v$.

Definition 3.1 G is a *Hadamard graph* (or simply "a graph") if

$G : \mathbf{B}_\delta^{E_x}(\mathbf{0}) \times \mathbf{B}_\delta^{F_x}(\mathbf{0}) \to \mathbf{B}_\delta^{E_x}(\mathbf{0})$,

$G(v_1, 0) = 0 \quad \forall\ v_1 \in \mathbf{B}_\delta^{E_x}(\mathbf{0})$ and

$\Phi : (v_1, v_2) \mapsto v_1 + v_2 + G(v_1, v_2) \in \mathbf{B}_{2\delta}^{T_xM}(\mathbf{0})$ is a C^1- diffeomorphism onto its image (see Figure 1).

The foliation $\mathcal{L}$ associated to the graph G is the foliation whose leaves are parametrized on $v_2 \in \mathbf{B}_\delta^{F_x}(\mathbf{0}) \subset F_x$, with constant v_1, by the diffeomorphism:

$$\exp_x(\Phi(v_1, \cdot)) = \exp_x(v_1 + \cdot + G(v_1, \cdot))$$

In Figure 1 we draw the foliation $\mathcal{L}$. To simplify the notation we do not write the exponential map $\exp_x$ and denote $y = v_1 + v_2 + G(v_1, v_2)$. The leaf containing y is written by $\mathcal{L}(y)$.




Figure 1: The foliation $\mathcal{L}$ associated to a Hadarmard graph (In this figure we do not write the exponential map $\exp_x$)

Definition 3.2 Dispersion of G

The dispersion of the graph G is

$$\operatorname{disp} G = \max_{v \in \mathbf{B}_\delta^{T_x M}(\mathbf{0})} \left\{ \left\| \frac{\partial G}{\partial v_2}(v) \right\| \right\},$$

where $v = v_1 + v_2$ and $\partial G / \partial v_2$ denotes the Fréchet derivative of

$$G(v_1, \cdot) \colon \mathbf{B}_\delta^{F_x}(\mathbf{0}) \to \mathbf{B}_\delta^{E_x}(\mathbf{0})$$

with a constant value of $v_1 \in \mathbf{B}_\delta^{E_x}(\mathbf{0})$.

We will denote by m^W the Lebesgue measure along an embedded local submanifold $W \subset M$.

Assertions 3.3

$$T_y \mathcal{L}(y) = \left(\operatorname{Id}|_{F_x} + \frac{\partial G}{\partial v_2}(v) \right) F_x, \tag{6}$$

$$m^{\mathcal{L}(y)}(\mathcal{L}(y)) \leq [(1 + \operatorname{disp} G)\delta]^{\dim F}, \tag{7}$$

For all $\varepsilon > 0$ *there exists* $c > 0$ *such that, if* $\operatorname{disp} G \leq c$, *then*

$$\operatorname{dist}(T_y \mathcal{L}(y), F_x) < \frac{\varepsilon}{2} \ \forall \ y \in \operatorname{Im}(\Phi).$$

For such a value of $c > 0$ (depending on $\varepsilon > 0$), there exists $\delta_1 > 0$ such that, if $\operatorname{dist}(x, y) < \delta_1$, *then* $y \in \operatorname{Im}(\Phi)$ *and*

$$\operatorname{dist}(T_y\mathcal{L}(y), F_y) < \varepsilon \tag{8}$$

Proof: The above assertions follow from the properties imposed in the definition of Hadamard graphs and their associated foliations, and from the definition of dispersion. In particular (8) holds because of the continuous dependence of the splitting $E_y \oplus F_y$ on the point y. □

3.4 Iterating the local foliation $\mathcal{L}$ by $\mathbf{f^n}$

Denote by $B^n_\delta(x)$ the dynamical ball defined as

$$B^n_\delta(x) := \{y \in M : \operatorname{dist}(f^j(x), f^j(y)) < \delta \ \forall\, 0 \le j \le n\}.$$

Take any graph G in $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$ such that $\operatorname{disp} G < 1/2$, and consider its associated local foliation $\mathcal{L}$. Construct the image $f^n(\mathcal{L})$ in the dynamical ball $B^n_\delta(x)$, i.e.:

$$f^n(\mathcal{L} \cap B^n_\delta(x)) := \exp^{-1}_{f^n(x)} f^n \exp_x(v_1 + v_2 + G(v_1, v_2))$$

for all $(v_1, v_2) \in \mathbf{B}_\delta^{E_x}(\mathbf{0}) \times \mathbf{B}_\delta^{F_x}(\mathbf{0})$ such that $\exp_x(v_1 + v_2 + G(v_1, v_2)) \in B^n_\delta(x)$.

Lemma 3.5 (Reformulation of Lemma 4 of [M81])
There exists $0 < c' < 1/2$ depending only on f, such that for all $0 < c < c'$ there exist $\delta_0, n_0 > 0$ such that for any point $x \in M$, if $\mathcal{L}$ is the local foliation associated to a graph G defined on T_xM with

$$\operatorname{disp} G < c,$$

then for all $n \ge 0$ the iterated foliation $f^n(\mathcal{L} \cap B^n_{\delta_0}(x))$ is contained in the associated foliation of a graph G_n defined on $T_{f^n(x)}M$, and

$$\operatorname{disp} G_n < c \text{ for all } n \ge n_0. \tag{9}$$

Besides, for all $y \in B^n_{\delta_0}(x)$ the image $f^n(\mathcal{L}(y) \cap B^n_{\delta_0}(x))$ is contained in a single leaf of the foliation associated to G_n.

Proof:
Step 1. Choose $n_0 \ge 0$ such that

$$\left\|df^n|_{E_z}\right\| \, \left\|df^{-n}|_{F_{f^n(z)}}\right\| < 1 \quad \forall\ n \ge n_0, \quad \forall\ x \in M. \tag{10}$$

For such a fixed value of n_0, take $\delta_0 > 0$ so that for all $x \in M$, for all $0 \le n \le n_0$, and for all the graphs G defined in $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$ with $\operatorname{disp} G < 1/2$, there exists a graph G_n defined on $\mathbf{B}_\delta^{T_{f^n(x)}M}(\mathbf{0})$ satisfying the following conditions:

$$\forall\ y = \exp_x(v_1 + v_2 + G(v_1, v_2)) \in B_{\delta_0}(x) \ \ \exists\ (u_1, u_2) \in \mathbf{B}_\delta^{E_{f^n(x)}}(\mathbf{0}) \times \mathbf{B}_\delta^{F_{f^n(x)}}(\mathbf{0}), \tag{11}$$

$$\text{where } u_1 \text{ depends only on } v_1 \tag{12}$$

$$\text{and } f^n(y) = \exp_{f^n(x)}(u_1+u_2+G_n(u_1,u_2)). \tag{13}$$

In the Appendix, Assertion 5.1, we show that such $\delta_0 > 0$ exists. We note that the condition above states $y \in B_{\delta_0}(x)$, which is not a dynamical ball, that the assertion is true for any initial graph G with dispersion smaller than $1/2$, and that Statements (11), (12) and (13) hold *only* for $0 \leq n \leq n_0$.

The condition that u_1 depends only on v_1 means that the images $f^n(z)$ for all the points $z \in B_{\delta_0}(x)$ that are in the same leaf $\mathcal{L}(y)$ associated to the graph G, are contained in *the same* leaf associated to the graph G_n.

Step 2. With $\delta_0 > 0$ fixed as above, we shall check the following statement:

There exists $0 < c' < 1/2$ such that for all the graphs G with $\operatorname{disp} G < c'$ and for all $n \geq 0$, if G_n is a graph defined in $\mathbf{B}_\delta^{T_{f^n(x)}M}(\mathbf{0})$ satisfying (11), (12) and (13), then:

$$\left\| \frac{\partial G_n}{\partial u_2}(u_1,u_2) \right\| \leq \|df^n|_{E_x}\| \cdot \operatorname{disp} G \cdot \left\|df^{-n}|_{F_{f^n(x)}}\right\| \ \ \forall\ y \in B^n_{\delta_0}(x) \tag{14}$$

We check this statement in Assertion 5.2 of the Appendix.

Step 3. Due to the construction of δ_0 in the step 1, Inequality (14) holds, in particular for $n = n_0$ for any G such that $\operatorname{disp} G < c'$. Therefore, using Inequalities (10) and (14), and Definition 3.2, we obtain:

$$\operatorname{disp} G_{n_0} \leq \operatorname{disp} G < c' \ \ \forall\ G \text{ s.t. } \operatorname{disp} G < c',$$

Moreover, if $\operatorname{disp} G < c < c'$, then $\operatorname{disp} G_{n_0} < c < c'$.

Step 4. From the construction of δ_0 in the step 1, using that $\operatorname{disp} G_{n_0} < c < c' < 1/2$ we deduce that there exists the graph G_n also for all $n_0 \leq n \leq 2n_0$. Besides $\|df^n|_{E_x}\| \cdot \|df^{-n}|_{F_{f^n(x)}}\| < 1$ for all $n \geq n_0$. So, applying Inequality (14), we deduce that $\operatorname{disp} G_n < c$ also for all $n_0 \leq n \leq 2n_0$. Thus, we can apply inductively the property of $\delta_0 > 0$ constructed in the Step 1, and Assertions (10), (11), (12), (13) and (14), to deduce that there exists the graph G_n for all $n \geq 0$ and $\operatorname{disp} G_n < c$ for all $n \geq n_0$. □

Once the constant c' of Lemma 3.5 is fixed, depending only on f, one obtains the following property that allows to move the reference point x (used to construct the graph G on $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$), preserving the *same* associated local foliation $\mathcal{L}$ and the uniformity of the upper bound of its dispersion:

Lemma 3.6 *For all $0 < c < c'$ there exists $\delta_1 > 0$ such that, for any $x \in M$ and for any graph G with $\operatorname{disp}(G) < c/2$ defined in $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$, the associated foliation $\mathcal{L}$ in the neighborhood $B_{\delta_1}(x)$ is also associated to a graph G' defined in $\mathbf{B}_\delta^{T_zM}(\mathbf{0})$, for any $z \in B_{\delta_1}(x)$. Besides: $\operatorname{disp}(G') < c$.*

Proof: The splitting $E_z \oplus F_z$ depends continuously on $z \in M$. Then, π_{E_z} and π_{F_z} also depend continuously on z. Therefore, for all $\epsilon > 0$ there exists $\delta_1 > 0$ such that

$$\|\pi_{F_x}|_{E_z}\| < \epsilon, \quad \|\pi_{E_x}|_{F_z}\| < \epsilon \quad \text{if } \operatorname{dist}(x,z) < \delta_1.$$

(For simplicity in the notation, in the above inequalities we have omitted the derivative of $\exp_x^{-1} \circ \exp_z$, which identifies T_zM with T_xM).

First, we claim that if $\delta_1 > 0$ is small enough, then for any graph G defined on $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$, and for any point z such that $\mathrm{dist}(z,x) < \delta_1$, there exists a graph G' defined on $\mathbf{B}_\delta^{T_zM}(\mathbf{0})$ such that the local foliations associated to G and G' coincide in an open set where both are defined. In fact, G' should satisfy the following equations:

$$u_1 + u_2 + G'(u_1,u_2) = v_1 + v_2 + G(v_1,v_2), \tag{15}$$

$$u_1, G'(u_1,u_2) \in E_z, \qquad u_2 \in F_z, \qquad G'(u_1,0) = 0.$$

Since by hypothesis G is a graph, it is C^1 and

$$v_1, \quad G(v_1,v_2) \in E_x, \quad v_2 \in F_x, \quad G(v_1,0) = 0.$$

The above equations are solved by

$$u_1 := \pi_{E_z}(v_1), \qquad u_2 := \pi_{F_z}(v_1 + v_2 + G(v_1,v_2)), \tag{16}$$

$$G' = -u_1 + \pi_{E_z}(v_1 + v_2 + G(v_1,v_2)). \tag{17}$$

The two equalities in (16) define a local diffeomorphism $\Psi : (v_1,v_2) \mapsto (u_1,u_2)$. In fact, on the one hand $u_1 = \pi_{E_z}|_{E_x}(v_1)$, where $\pi_{E_z}|_{E_x}$ is a diffeomorphism (which is linear and uniformly near the identity map, independently of the graph G). On the other hand, for v_1 constant, the derivative with respect to v_2 of $\pi_{F_z}(v_1+v_2+G(v_1,v_2))$ is $\pi_{F_z}|_{T_y\mathcal{L}(y)}$, which is, independently of the graph G, uniformly near $\pi_{F_x}|_{T_y\mathcal{L}(y))} = Id|_{F_x}$. Thus, Ψ is a local diffeomorphism, which, besides, is C^1 near the identity map, if δ_1 is chosen small enough (independently of the given graph G).

From the above construction, we deduce that the composition of the mapping $\Psi : (u_1,u_2) \mapsto (v_1,v_2)$ with the mapping $(v_1,v_2) \mapsto G'$ defined by (17), is of C^1 class. Therefore $G'(u_1,u_2)$ depends C^1 on (u_1,u_2). Besides $G'(u_1,0) = 0$ because $G(v_1,0) = 0$. Due to Identity (15), the application ϕ' defined by $\Phi'(u_1,u_2) := u_1 + u_2 + G'(u_1,u_2)$ coincides with the application $\Phi(v_1,v_2) := v_1 + v_2 + G(v_1,v_2)$. This latter mapping Φ, due to Definition 3.1, is a local diffeomorphism. So Φ' is also a local diffeomorphism and G' satisfies Definition 3.1 to be a Hadamard graph. The first claim is proved.

The diffeomorphism $\Psi : (v_1,v_2) \mapsto (u_1,u_2)$ as constructed above, converges to the identity map in the C^1 topology, when $\delta_1 \to 0^+$, and uniformly for all the graphs G defined in $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$. Thus, by Identity (15), $\|G' - G\|_{C^1}$ converges uniformly to zero, independently of the given graph G, when $\delta_1 \to 0$. This implies, in particular, that $\partial G'(u_1,u_2)/\partial u_2$ converges uniformly to $\partial G(v_1,v_2)/\partial v_2$ when $\delta_1 \to 0$. Thus, for any constant $c/2 > 0$ there exists $\delta_1 > 0$, which is independent of the graph G, such that $|\,\mathrm{disp}(G') - \mathrm{disp}(G)| < c/2$. In other words, $\mathrm{disp}(G') < c$ for all G with $\mathrm{disp}(G) < c/2$, as wanted. □

We are ready to prove the following Proposition, for all $f \in \mathrm{Diff}^1(M)$ with a dominated splitting $TM = E \oplus F$.

Proposition 3.7 *For all $\varepsilon > 0$ there exist $\delta_0, K, n_0 > 0$, and a finite family of local foliations $\mathcal{L}$, each one defined in an open ball of a given finite covering of M with δ_0-balls, such that:*
(a) *$\mathcal{L}$ is C^1- trivializable and its leaves are* $\dim F$*-dimensional*
(b) $\operatorname{dist}\left(F_{f^n(x)}, T_{f^n(x)} f^n(\mathcal{L}(x))\right) < \varepsilon \ \ \forall\ x,\ \forall\ n \geq n_0$
(c) *$\forall\ n \geq 0$ and $\forall\ x, y$ such that $y \in B^n_{\delta_0}(x)$:*

$$m^{f^n(\mathcal{L}(y))}(f^n(\mathcal{L}(y) \cap B^n_{\delta_0}(x))) \leq K$$

(d) *$\forall\ n \geq 0$ and $\forall\ x \in M$:*

$$e^{-n\varepsilon} K^{-1} \leq \frac{\left|\det df^n_x|_{T_x(\mathcal{L}(x))}\right|}{|\det df^n_x|_{F_x}|} \leq K e^{n\varepsilon}.$$

Proof: Consider the constant c' that is determined by Lemma 3.5. For each point $x \in M$ construct a local foliation $\mathcal{L}$ from a graph G defined on T_xM, with dispersion smaller than a constant $c/2$ such that $0 < c < c' < 1/2$. The constant c will be fixed later taking into account the given value of $\varepsilon > 0$. After Lemma 3.6, there exists $\delta_1 > 0$ such that, for all $x \in M$ the graph G defined on $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$ is redefined on $\mathbf{B}_\delta^{T_zM}(\mathbf{0})$, for any point $z \in B_{\delta_1}(x)$, preserving the same associated foliation and having dispersion upper bounded by c. Fix δ_0, n_0 (depending on c) by Lemma 3.5 and such that $\delta_0 < \delta_1$. For any given finite covering of M with balls $B_{\delta_0}(x_i)$, fix a finite family $\{\mathcal{L}_i\}_{1 \leq i \leq k}$ of local foliations so constructed, one in each ball of the covering.

By the definition of graph, each foliation $\mathcal{L}$ of the finite family constructed above, is C^1-trivializable and its leaves are dim F-dimensional. Thus Assertion **(a)** holds.

From Inequality (8), given $\varepsilon' > 0$ (a fixed value of $\varepsilon' > 0$ will be determined later), there exists $c > 0$ such that, if $\operatorname{disp}(G) < c$ then

$$\operatorname{dist}(T_x(\mathcal{L}(x), F_x)) < \varepsilon' \ \ \forall\ x \in M. \tag{18}$$

Recall that δ_0, n_0 (depending on c, which depends on ε') were defined by Lemma 3.5. Therefore, each leaf $f^n(\mathcal{L}(y) \bigcap B^n_{\delta_0}(x))$ is part of a single leaf of a foliation associated to a graph G_n, for all $n \geq 0$. Besides, Lemma 3.5 states that

$$\operatorname{disp} G_n < c \ \ \forall\, n \geq n_0 \tag{19}$$

From Inequalities (7) and (19) we deduce that

$$m^{f^n(\mathcal{L}(y))}(f^n(\mathcal{L}(y) \cap B^n_{\delta_0}(x))) \leq m^{f^n(\mathcal{L}(y))}(f^n(\mathcal{L}(y)) \leq$$

$$[(1 + \operatorname{disp} G_n)\, \delta\,]^{\dim F} < [(1 + c)\, \delta\,]^{\dim F} \quad \forall\ n \geq n_0.$$

Thus, there exists $K > 0$ such that

$$m^{f^n(\mathcal{L}(y))}(f^n(\mathcal{L}(y) \cap B^n_{\delta_0}(x))) \leq K \quad \forall\ n \geq 0.$$

So, Assertion **(c)** of Proposition 3.7 is proved for each fixed value of $\varepsilon' > 0$.

Next, we prove **(d)**. From Inequalities (18) and (19), we deduce that

$$\operatorname{dist}\left(F_{f^n(x)}, T_{f^n(x)}f^n(\mathcal{L}(x))\right) < \varepsilon' \ \ \forall\ x \in M,\ \forall\ n \geq n_0. \tag{20}$$

Finally, we fix $\varepsilon' > 0$ (depending on the given value of $\varepsilon > 0$), such that $0 < \varepsilon' < \varepsilon$ and such that for all the $\dim F$-dimensional sub-bundles L that satisfy $\operatorname{dist}(L, F) < \varepsilon'$, the following inequality holds:

$$e^{-\varepsilon} \leq \frac{\left|\det df_x|_{L(x)}\right|}{\left|\det df_x|_{F_x}\right|} \leq e^{\varepsilon} \ \ \forall\ x \in M. \tag{21}$$

Therefore, (20) implies:

$$e^{-\varepsilon} \leq \frac{\left|\det df_{f^j(x)}|_{T_{f^j(x)}f^j\mathcal{L}(x)}\right|}{\left|\det df_{f^j(x)}|_{F_{f^j(x)}}\right|} \leq e^{\varepsilon} \ \ \forall\ x \in M, \ \ \forall\ j \geq n_0.$$

The latter inequality implies **(d)**. Finally, **(b)** is obtained from (20) taking into account that ε' was chosen smaller than ε. □

End of the proof of Lemma 2.1:

For the given value of $\varepsilon > 0$, we construct $\delta_0, K > 0$ as in Proposition 3.7. Consider any finite partition $\alpha = \{A_h\}_{1 \leq h \leq k}$, where $k = \#(\alpha)$, such that $\operatorname{diam}\alpha = \max_{h=1\ldots k} \operatorname{diam}\{A_h\} < \delta_0$.

For each $A \in \alpha$ construct an open set $V_A \subset M$ also of diameter smaller than δ_0, containing A. Construct a $\dim F$ local foliation $\mathcal{L}_A$ in V_A satisfying Proposition 3.7. Construct also a C^1 submanifold W_A transversal to $\mathcal{L}_A$.

Take $\alpha^n = \vee_{j=0}^{n} f^{-j}(\alpha) = \{X_i\}_{1 \leq i \leq k_n}$, where $k_n = \#(\alpha^n)$. For all $X_i \in \alpha^n$, there exists $A_{h_i} \in \alpha$ such that $X_i \subset A_{h_i}$. Denote $\mathcal{L}_i := \mathcal{L}_{A_{h_i}}$ and $W_i := W_{A_{h_i}}$. Since $\mathcal{L}_i$ is C^1-trivializable, by Fubini's Theorem we have:

$$m(C) = \sum_{i=1}^{k_n} \int_{z \in W_i} d\mu^{W_i} \int_{y \in \mathcal{L}_i(z)} \mathbf{1}_{C \bigcap X_i}\, \phi_i\, dm^{\mathcal{L}_i(z)} \qquad \forall\ C \in \mathcal{B}, \tag{22}$$

where $\mathcal{B}$ is the Borel sigma-algebra, $\mathbf{1}_{C \bigcap X_i}$ is the characteristic function of the set $C \bigcap X_i$, and ϕ_i is a continuous function which depends on $A_{h_i} \in \alpha$. Precisely, ϕ_i is the Jacobian of the C^1-trivialization of the foliation $\mathcal{L}_i$. So, there exist at most $k = \#(\alpha)$ different local foliations $\mathcal{L}_i$, k different continuous functions ϕ_i, and k different transversal manifolds W_i, which allow Formula (22) work for any value of n and for any $C \in \mathcal{B}$.

Denote $\widehat{y} = f^n(y) \in f^n(\mathcal{L}_i(z) \bigcap X_i) =: \mathcal{L}_i^n(z)$:

$$m(C) = \sum_{i=1}^{k_n} \int_{z \in W_i} d\mu^{W_i} \int_{\widehat{y} \in \mathcal{L}_i^n(z)} [\mathbf{1}_{C \cap X_i}\ \phi_i](f^{-n}(\widehat{y}))\left|\det df^{-n}|_{T_{\widehat{y}}\mathcal{L}_i^n}\right| dm^{\mathcal{L}_i^n(z)}. \tag{23}$$

By Part **(d)** of Proposition 3.7:

$$\left|\det df^{-n}|_{T_{\widehat{y}}\mathcal{L}_i^n}\right| \leq K e^{n\varepsilon}\ \left|\det df^{-n}|_{F_{\widehat{y}}}\right|. \tag{24}$$

Recall Formula (3) defining $\psi_n(y)$. Since $f^n(y) = \widehat{y}$, we have

$$\left|\det df^{-n}|_{F_{\widehat{y}}}\right| = \psi_n(f^{-n}(\widehat{y})).$$

Joining with Inequality (24) and Equality (23), we obtain:

$$\left|\det df^{-n}|_{T_{\widehat{y}}\mathcal{L}_i^n}\right| \leq Ke^{n\varepsilon}e^{\psi_n(f^{-n}(\widehat{y}))} \tag{25}$$

$$m(C) \leq Ke^{n\varepsilon}\sum_{i=1}^{k_n}\int_{z\in W_i} d\mu^{W_i}\int_{\widehat{y}\in\mathcal{L}_i^n(z)}[\mathbf{1}_{C\cap X_i}\ \phi_i](f^{-n}(\widehat{y}))\, e^{\psi_n(f^{-n}(\widehat{y}))}\, dm^{\mathcal{L}_i^n(z)}. \tag{26}$$

By Radon-Nykodim Theorem, construct the finite measure ν_n satisfying:

$$\int h\, d\nu_n = \sum_{i=1}^{k_n}\int_{z\in W_i} d\mu^{W_i}\int_{\widehat{y}\in\mathcal{L}_i^n(z)}[(\mathbf{1}_{X_i}\cdot\phi_i\cdot h)\circ f^{-n}]\ (\widehat{y})\ dm^{\mathcal{L}_i^n(z)}\ \ \forall\, h\in C^0(M,\mathbb{R}).$$

From Inequality (26) and the above definition of ν_n, we obtain:

$$m(C) \leq Ke^{n\varepsilon}\int \mathbf{1}_C\, e^{\psi_n}\, d\nu_n \leq Ke^{n\varepsilon}\int_C e^{\psi_n}\, d\nu_n.$$

Statement (ii) of Lemma 2.1 is proved. Let us prove Statement (i). We must show that there exists a constant $K_0 > 0$, independent of n, such that $\nu_n(X) \leq K_0$ for all $X \in \alpha^n$, and for all $n \geq 0$. In fact, recall that $\mathcal{L}_i^n(z) = f^n(\mathcal{L}_i(z)\cap X_i) \subset f^n(\mathcal{L}_i(z)\cap B_{\delta_0}(y))$ for all $z \in W_i$ and for all $y \in \mathcal{L}_i(z)\cap X_i$. Thus, applying Property **(c)** of Proposition 3.7 we have

$$m^{\mathcal{L}_i^n}(\mathcal{L}_i^n(z)) \leq K_1,$$

for some constant $K_1 > 0$ which is independent of n. From the construction of the measure ν_n:

$$\nu_n(X_i) = \int_{z\in W_i} d\mu^{W_i}\int_{\widehat{y}\in\mathcal{L}_i^n(z)}[(\mathbf{1}_{X_i}\cdot\phi_i)\circ f^{-n}]\,(\widehat{y})\, dm^{\mathcal{L}_i^n(z)} \leq$$

$$\mu^{W_i}(W_i)\,\|\phi_i\|_{C^0}\ m^{\mathcal{L}_i^n}(\mathcal{L}_i^n(z)) \leq K_1\,\mu^{W_i}(W_i)\,\|\phi_i\|_{C^0}.$$

Since the number of different local foliations $\mathcal{L}_i$ is equal to the number k of pieces of the given partition α, which is independent of n, we obtain:

$$\nu_n(X_i) \leq K_1 \max_{A\in\alpha}\{\mu^{W_A}(W_A)\,\|\phi_A\|_{C^0}\} =: K_0,$$

where K_0 is a constant that depends only on the partition α and not on n. $\square$

4 Proof of Lemma 2.2

In Lemma 2.2, a measure $\mu \in \mathcal{P}_f$ and two small numbers $\varepsilon > 0$ and $\delta > 0$, are arbitrarily given. We must construct an adequate finite partition α of M, with diameter smaller than δ, and a sequence $\{C_n\}_{n\geq 0} \subset \mathcal{B}$, satisfying the statements **(iii)** and **(iv)** of Lemma 2.2.

Assume that dist* and $\varepsilon^* > 0$ are already chosen. We need to construct $\{C_n\}$ such that the basin $B_{\varepsilon^*}(\mu)$ of statistical ε^*-weak attraction of μ is contained in the set of points $x \in M$ satisfying $x \in C_n$ for infinitely values of $n \in \mathbb{N}$. Using Definition 1.2 of $B_{\varepsilon^*}(\mu)$, we choose:

$$C_n = \{x \in M : \text{dist}^*(\sigma_{n,x}, \mu) < \varepsilon^*\}, \tag{27}$$

where $\sigma_{n,x}$ is the empirical probability of the future orbit of x (see Definition 1.1). So, statement **(iii)** of Lemma 2.2 is satisfied.

Let us construct the partition α to prove statement **(iv)**. Take $\delta_1 > 0$ such that $\text{dist}(x, y) < \delta_1 \Rightarrow |\psi(x) - \psi(y)| < \varepsilon/5$.
Take α such that $\text{diam}(\alpha) \leq \min(\delta, \delta_1)$, $\mu(\partial X) = 0 \ \forall \ X \in \alpha$. This construction implies

$$\lim_{n\to+\infty} \mu_n(X) = \mu(X) \tag{28}$$

for all $X \in \alpha^q = \bigvee_{j=0}^{q} f^{-1}(\alpha)$, for all $q \in \mathbb{N}$ and for all $\{\mu_n\}_n \subset \mathcal{P}$ such that $\lim_n^* \mu_n = \mu$. Also

$$|\psi_n(y) - \psi_n(x)| \leq \sum_{j=0}^{n-1} |\psi(f^j(y)) - \psi(f^j(x))| \leq \frac{n\varepsilon}{5} \ \ \forall \ x, y \in X, \forall X \in \alpha^n. \tag{29}$$

Recall that $h_\mu(\alpha) := \lim_{q\to+\infty} H(\alpha^q, \mu)/q$, where $H(\alpha^q, \mu) := -\sum_{X\in\alpha^q} \mu(X) \log \mu(X)$.
Fix $q \in \mathbb{N}^+$ s.t. $H(\alpha^q, \mu)/q < h_\mu(\alpha) + \varepsilon/5$. From (28):

$$\lim_{n\to+\infty} \frac{H(\alpha^q, \mu_n)}{q} = \frac{H(\alpha^q, \mu)}{q} < h_\mu(\alpha) + \frac{\varepsilon}{5} \tag{30}$$

for any sequence $\mu_n \in \mathcal{P}$ such that $\lim_n^* \mu_n = \mu$.

Choose $\{\varphi_i\}_{i\geq 1}$ dense in $C^0(M, [0, 1])$ and define dist* in $\mathcal{P}$:

$$\text{dist}^*(\mu_1, \mu_2) := \left| \int \psi \, d\mu_1 - \int \psi \, d\mu_2 \right| + \sum_{i=1}^{\infty} \frac{\left| \int \varphi_i \, d\mu_1 - \int \varphi_i \, d\mu_2 \right|}{2^i}. \tag{31}$$

Using (30), fix $0 < \varepsilon^* < \varepsilon/5$ such that

$$\sigma \in \mathcal{P}, \ \text{dist}^*(\sigma, \mu) \leq \varepsilon^* \Rightarrow |H(\alpha^q, \sigma) - H(\alpha^q, \mu)| \leq \frac{q\varepsilon}{5}. \tag{32}$$

Such a value of ε^* exists because if not, we would construct a sequence of probability measures μ_n, converging to μ and such that $|H(\alpha^q, \mu_n) - H(\alpha^q, \mu)| > q\,\varepsilon/5$ for all $n \in \mathbb{N}$. This inequality contradicts the equality at left in (30).

For each fixed $n \geq 0$, consider $\alpha^n \bigvee \{C_n\} := \{X_i \cap C_n : \ \ X_i \in \alpha^n, \ \ X_i \cap C_n \neq \emptyset\}$. Denote $k_n := \#(\alpha \bigvee \{C_n\})$. For each $C_n \cap X_i \in \alpha \bigvee \{C_n\}$, choose one and only one

point $x_i \in C_n \bigcap X_i$. Construct the integral appearing in the left member of the inequality **(iv)** of Lemma 2.2, and apply Inequality (29):

$$I_n := \int e^{\psi_n}\, d\nu_n = \sum_{i=1}^{k_n} \int_{y \in C_n \cap X_i} e^{\psi_n(y)} d\nu_n(y) \leq \sum_{i=1}^{k_n} e^{n\varepsilon/5} e^{\psi_n(x_i)} \nu_n(C_n \cap X_i). \quad (33)$$

By hypothesis $\nu_n(X) \leq K$ for all $X \in \alpha^n$. So:

$$I_n \leq K e^{n\varepsilon/5} \sum_{i=1}^{k_n} e^{\psi_n(x_i)} \quad (34)$$

Define $p_i := e^{\psi_n(x_i)}/L$, where $L := \sum_{i=1}^{k_n} e^{\psi_n(x_i)}$. Note that $\sum_{i=1}^{k_n} p_i = 1$. Apply the "equality" condition in the Inequality of Jensen:

$$\log \sum_{i=1}^{k_n} e^{\psi_n(x_i)} = \sum_{i=1}^{k_n} \psi_n(x_i) p_i - \sum_{i=1}^{k_n} p_i \log p_i. \quad (35)$$

Taking logarithm in (34) and using (35), we obtain:

$$\log I_n \leq \log K + \frac{n\epsilon}{5} + \sum_{i=1}^{k_n} \psi_n(x_i) p_i - \sum_{i=1}^{k_n} p_i \log p_i.$$

From Equalities (2) and (3):

$$\sum_{i=1}^{k_n} \psi_n(x_i) p_i = \sum_{i=1}^{k_n} \sum_{j=0}^{n-1} \int p_i\, \psi\ d\delta_{f^j(x_i)},$$

and thus:

$$\log I_n \leq \log K + \frac{n\varepsilon}{5} + \sum_{i=1}^{k_n} \sum_{j=0}^{n-1} \int p_i\, \psi\ d\delta_{f^j(x_i)} - \sum_{i=1}^{k_n} p_i \log p_i. \quad (36)$$

Recall Definition 1.1 of the empirical probabilities $\sigma_{n,x}$. We construct $\mu_n \in \mathcal{P}$ by

$$\mu_n := \frac{1}{n} \sum_{i=1}^{k_n} \sum_{j=0}^{n-1} p_i\ \delta_{f^j(x_i)} = \sum_{i=1}^{k_n} p_i \sigma_{n,x_i}. \quad (37)$$

Since $x_i \in C_n$ we have that $\mathrm{dist}^*(\sigma_{n,x_i}, \mu) < \varepsilon^*$ (see Equality (27)). Since the ε^*-balls defined with the metric dist^* by Equality (31) are convex, and μ_n is a convex combination of the measures σ_{n,x_i}, we deduce

$$\mathrm{dist}^*(\sigma_{n,x_i}, \mu) \leq \varepsilon^* \Rightarrow \mathrm{dist}^*(\mu_n, \mu) \leq \varepsilon^*.$$

From the construction of dist^* by Equality (31), we obtain $\left|\int \psi\, d\mu_n - \int \psi\, d\mu\right| \leq \varepsilon^* < \varepsilon/5$. Therefore:

$$\int \psi\, d\mu_n \leq \int \psi\, d\mu + \frac{\varepsilon}{5},$$

which joint with (36) and (37) implies:

$$\log I_n \leq \log K + \frac{2n\varepsilon}{5} + n\int \psi\, d\mu - \sum_{i=1}^{k_n} p_i \log p_i.$$

In Assertion 5.3 of the Appendix, we check the following statement:

There exists $n_0 \geq 0$ *such that*

$$-\sum_{i=1}^{k_n} p_i \log p_i \leq \frac{n\,\varepsilon}{5} + \frac{n}{q} H(\alpha^q, \mu_n) \quad \forall n \geq n_0$$

Therefore:

$$\log I_n \leq \log K + \frac{3n\varepsilon}{5} + n\int \psi\, d\mu + \frac{nH(\alpha^q, \mu_n)}{q}.$$

By construction of the positive number ε^* in (32) and since $\text{dist}^*(\mu_n, \mu) < \varepsilon^*$, we have

$$\left| \frac{H(\alpha^q, \mu_n)}{q} - \frac{H(\alpha^q, \mu)}{q} \right| \leq \frac{\varepsilon}{5}.$$

So

$$\log I_n \leq \log K + \frac{4n\varepsilon}{5} + n\int \psi\, d\mu + \frac{n\,H(\alpha^q, \mu)}{q}.$$

Using the choice of q by Inequality (30) at the beginning of the proof, we conclude:

$$\log I_n \leq \log K + n\,\varepsilon + n\int \psi\, d\mu + n\, h_\mu(\alpha),$$

which shows the Assertion **(iv)**, ending the proof of Lemma 2.2. □

5 Appendix

In this section we check some technical assertions that were used in the proofs of Sections 3 and 4.

Assertion 5.1 *Let* $\delta > 0$ *be such that for all* $x \in M$

$$\exp_x : \{v \in T_xM : \|v\| \leq 3\delta\} \to B_{3\delta}(x) \subset M$$

is a diffeomorphism. Let $n_0 > 0$.

Then, there exists $0 < \delta_0 < \delta$ *such that* $\forall\ x \in M,\ \forall\ 0 \leq n \leq n_0$ *and for all the graphs* G (defined in $\mathbf{B}_\delta^{T_xM}(\mathbf{0}) \subset T_xM$) *such that*

$$\text{disp } G < 1/2,$$

there exists a graph G_n (defined in $\mathbf{B}_\delta^{T_{f^n(x)}M}(\mathbf{0})$) *satisfying:*

$$\forall\ y = \exp_x(v_1 + v_2 + G(v_1, v_2)) \in B_{\delta_0}(x)\ \exists\ (u_1, u_2) \in \mathbf{B}_\delta^{E_{f^n(x)}}(\mathbf{0}) \times \mathbf{B}_\delta^{F_{f^n(x)}}(\mathbf{0}) \quad (38)$$

$$u_1 \textit{ depends only on } v_1, \textit{ and} \quad (39)$$

$$f^n(y) = \exp_{f^n(x)}(u_1 + u_2 + G_n(u_1, u_2)). \quad (40)$$




Figure 2: The image $f(\mathcal{L}(y))$ of the leaves $\mathcal{L}(y)$ near x associated to the graph G, are associated to the graph G_1. (We do not write the exponential maps $\exp_x$, $\exp_{f(x)}$).

Proof: We will argue by induction on $n \in \mathbb{N}$, to show that for each $n \geq 1$, there exists $\delta_n > 0$ and G_n satisfying statements (38), (39), (40). To prove Assertion 5.1 it is enough to take $\delta_0 := \min\{\delta_1, \ldots, \delta_{n_0}\}$.

To simplify the notation along the proof, we will not write the exponential maps. From Definition 3.1, recall the construction of the diffeomorphism Φ obtained from the graph G, which is a trivialization of the associated local foliation $\mathcal{L}$ (See the upper frame of Figure 2). Precisely, each leaf $\mathcal{L}(v_1)$ is obtained for constant $v_1 \in \mathbf{B}_\delta^{E_x}(\mathbf{0})$, and parametrized by $v_2 \in \mathbf{B}_\delta^{F_x}(\mathbf{0})$ by the formula

$$\mathcal{L}(v_1) : v_2 \mapsto \Phi(v_1, v_2) := v_1 + v_2 + G(v_1, v_2), \quad \text{where } G(v_1, v_2) \in \mathbf{B}_\delta^{E_x}(\mathbf{0}).$$

Since $G(v_1,0)=0$ we have $v_1=\Phi(v_1,0)$. i.e. $v_1\in\mathcal{L}(v_1)\cap\mathbf{B}_\delta^{E_x}(\mathbf{0})$. Moreover, $\pi_{F_x}(v_1+v_2+G(v_1,v_2))=v_2$ for all $v_2\in\mathbf{B}_\delta^{F_x}(\mathbf{0})$. So

$$\pi_{F_x}\mathcal{L}(v_1)=\mathbf{B}_\delta^{F_x}(\mathbf{0}).$$

Besides, $\mathcal{L}(v_1)$ is uniformly transversal to E_x, for all G with $\mathrm{disp}(G)<1/2$. In fact, $T_{v_1}\mathcal{L}(v_1)=\mathrm{Im}(\mathrm{Id}|_{F_x}+\partial G/\partial v_2)$, the subspace F_x is transversal to E_x, and $\|\partial G/\partial v_2\|\leq \mathrm{disp}\,G<1/2$. Since the leaf $\mathcal{L}(x)$ intersects E_x at $\mathbf{0}$, we deduce that there exists $0<\delta'\leq\delta$, which is uniform for any G with $\mathrm{disp}(G)<1/2$, such that, if $\mathrm{dist}(x,y)<\delta'$ then y' belongs to some leaf of the foliation $\mathcal{L}$. In other words, $B_{\delta'}(x)\subset\mathrm{Im}(\Phi)$, i.e. there exists $(v_1,v_2)\in E_x\times F_x$ such that

$$\|v_1\|<\delta,\quad \|v_2\|<\delta,\quad y=\Phi(v_1,v_2)=v_1+v_2+G(v_1,v_2)\quad \text{if } y\in B_{\delta'}(x).$$

Recall that, by Definition 3.1, Φ is a diffeomorphism onto its image. Thus, for all $y\in B_{\delta'}(x)$, the point $\Phi^{-1}(y)=(v_1,v_2)\in\mathbf{B}_\delta^{E_x}(\mathbf{0})\times\mathbf{B}_\delta^{F_x}(\mathbf{0})$ depends C^1 on y. We take $0<\delta_1<\delta'$ such that if $y\in B_{\delta_1}(x)$, then

$$\|\pi_{E_{f(x)}}f(y)\|<\delta/2,\quad \|\pi_{F_{f(x)}}f(y)\|<\delta/2. \tag{41}$$

Such a value of $\delta_1>0$ exists, and is independent of the graph G, because f, π_{E_x} and π_{F_x} are uniformly continuous. Taking, if necessary, a smaller value of δ_1, the following two properties (A) and (B) are obtained for any graph G with $\mathrm{disp}(G)<1/2$ and for any $y\in B_{\delta_1}(x)$:

(A) The leaf $f(\mathcal{L}(y))$ intersects $\mathbf{B}_{\delta/2}^{E_{f(x)}}(\mathbf{0})\subset E_{f(x)}$ in a point u_1 (see Figure 2). In other words

$$\exists\ \ u_1\in E_{f(x)},\quad \|u_1\|<\delta/2,\quad f^{-1}(u_1)\in\mathcal{L}(y). \tag{42}$$

(B) The application $v_1\in E_x\ \mapsto\ u_1\in E_{f(x)}$ defined by (A) for all $y=\Phi(v_1,v_2)\in B_{\delta_0}(x)$, is independent of v_2, and is a diffeomorphism onto its image.

Property (A) is achieved due to the Implicit Function Theorem, since f is a diffeomorphism, $\mathrm{Im}\,df_x|_{E_x}=E_{f(x)}$, and the local foliation $\mathcal{L}$ is uniformly transversal to $\mathbf{B}_\delta^{E_x}(\mathbf{0})\subset E_x$, while its leaf $\mathcal{L}(x)$ intersects E_x at 0. Property (B) is obtained because f is a diffeomorphism and the mapping $f(v_1)\in f(E_x)\mapsto u_1\in E_{f(x)}$ is the holonomy along the leaves of the foliation $f(\mathcal{L})$, which is C^1 trivializable and uniformly transversal to both $f(E_x)$ and $E_{f(x)}$ (see Figure 2).

Let us show that the graph G_1 exists in $T_{f(x)}M$, satisfying Definition 3.1 and Assertions (38) (39) and (40) for all $y\in B_{\delta_1}(x)$. We write $y=\Phi(v_1,v_2)=v_1+v_2+G(v_1,v_2)$. We have already determined $u_1\in E_{f(x)}$ as a diffeomorphic function of v_1, independently of v_2. Let us determine $u_2\in F_{f(x)}$ and $G_1(u_1,u_2)\in E_{f(x)}$ such that $f(y)=u_1+u_2+G_1(u_1,u_2)$ (see Figure 2). Consider the equation:

$$f(v_1+v_2+G(v_1,v_2))=u_1+u_2+G_1(u_1,u_2)\ \ u_2\in F_{f(x)},\ \ G_1(u_1,u_2)\in E_{f(x)}, \tag{43}$$

which is solved by

$$u_2:=\pi_{F_{f(x)}}f(v_1+v_2+G(v_1,v_2))\in F_{f(x)}, \tag{44}$$

$$G_1(u_1,u_2) := -u_1 + \pi_{E_{f(x)}} f(v_1+v_2+G(v_1,v_2)) \in E_{f(x)}. \tag{45}$$

The application Ψ defined by $\Psi(v_1,v_2) \mapsto (u_1,u_2)$ where u_1 and u_2 are constructed as above, is of C^1 class. In fact, u_1 depends only on v_1, the mapping $v_1 \mapsto u_1$ is a diffeomorphism onto its image, and u_2 is constructed by Formula (44). Moreover, Ψ is C^1 invertible. In fact, on the one hand, we have that the application $v_1 \mapsto u_1$ is C^1 invertible and independent of v_2. On the other hand, for constant v_1 let us show that Formula (44) applies $v_2 \mapsto u_2$ C^1-diffeomorphically. Precisely, $G(v_1,0)=0$, $v_2 \in F_x$ $u_2 \in F_{f(x)}$, $G(v_1,v_2) \in E_x$ and $df_x : F_x \mapsto F_{f(x)}$ is invertible. Thus, $\pi|_{F_{f(x)}}\, df = df|_{F_x}\, \pi_{F_x}$. Taking derivatives in Equality (44) with respect to v_2 with constant v_1, we obtain:

$$\frac{\partial u_2}{\partial v_2} = df|_{F_x} \cdot \pi_{F_x} \left(Id|_{F_x} + \frac{\partial G(v_1,v_2)}{\partial v_2} \right) = df|_{F_x} = (df^{-1}|_{F_{f(x)}})^{-1}.$$

The second equality is due to $G(v_1,v_2) \in E_x$ for all (v_1,v_2), and so, the projection by π_{F_x} composed with any derivative of G is zero. We have proved that $\partial u_2/\partial v_2$ is invertible, and besides

$$\left(\frac{\partial u_2}{\partial v_2}\right)^{-1} = \frac{\partial v_2}{\partial u_2} = df^{-1}|_{F_{f(x)}} \tag{46}$$

We have shown that the application Ψ is a diffeomorphism onto its image. Now, we define the mapping Φ_1 by

$$\Phi_1 : (u_1,u_2) \mapsto u_1 + u_2 + G_1(u_1,u_2).$$

Φ_1 is a C^1 diffeomorphism onto its image, because its inverse is $\Psi \circ \Phi \circ f^{-1}$. So G_1 is C^1, and Φ_1 is the C^1 trivialization of its associated foliation, which is, by construction, $f(\mathcal{L})$.

Finally, (41) , (42) and (45) imply

$$\|G_1\| \leq \|u_1\| + \|\pi_{E_{f(x)}} f(y)\| < \delta/2 + \delta/2 = \delta \quad \text{and}$$

$$\Phi_1^{-1} f(B_{\delta_1}(x)) \subset \mathbf{B}_{\delta/2}^{E_{f(x)}}(\mathbf{0}) \times \mathbf{B}_{\delta/2}^{F_{f(x)}}(\mathbf{0}).$$

Thus, $G_1 : \Phi_1^{-1}(f(B_{\delta_1}(y))) \to \mathbf{B}_{\delta}^{E_{f(x)}}(\mathbf{0})$ can be C^1 extended to be a graph $G_1 : \mathbf{B}_{\delta}^{E_{f(x)}}(\mathbf{0}) \times \mathbf{B}_{\delta}^{F_{f(x)}}(\mathbf{0}) \to \mathbf{B}_{\delta}^{E_{f(x)}}(\mathbf{0})$.

We have completed the first step of the inductive proof, since we have proved the existence of $\delta_1 > 0$ and of the graph G_1 satisfying (38), (39) and (40). Naturally, disp G_1 is not necessarily upper bounded by $1/2$. So, we can not repeat exactly the same argument to prove the inductive step. Instead, we will prove that there exists an uniform constant $c_1 > 0$ such that

$$\operatorname{disp} G_1 < c_1 \quad \forall\ G \text{ such that } \operatorname{disp} G < 1/2. \tag{47}$$

If we prove Inequality (47) for some constant c_1, then we could end the inductive proof as follows:

Assume that for some $n \geq 0$ there exist $\delta_n, c_n > 0$ and a graph G_n defined in $\mathbf{B}_\delta^{T_{f^n(x)}M}(\mathbf{0})$ satisfying (38), (39) and (40) for all $y \in B_{\delta_n}(x)$, and such that $\operatorname{disp} G_n \leq c_n \ \ \forall\, G$ with $\operatorname{disp} G < 1/2$. Thus, we can repeat the above proof, putting $\min(\delta_n, \delta), c_n$ and G_n in the roles of $\delta, 1/2$ and G, respectively. We deduce that there exist $\delta_{n+1}, c_{n+1} > 0$ and a graph $G_{n+1} := (G_n)_1$, defined in $\mathbf{B}_\delta^{T_{f^{n+1}(x)}M}(\mathbf{0})$, which satisfies (38), (39) and (40) for all $y \in B_{\delta_{n+1}}(x)$, and such that $\operatorname{disp} G_{n+1} < c_{n+1}$ for all G for which $\operatorname{disp} G_n < c_n$. Thus, G_{n+1} satisfies (38), (39) and (40) for all G such that $\operatorname{disp} G < 1/2$. Therefore, the inductive proof will be completed, once we show Inequality (47).

So, let us find a constant c_1 satisfying Inequality (47). To find c_1, we upper bound $\|\partial G_1(u_1,u_2)/\partial u_2\|$. From (45) and taking into account that $\pi_{E_{f(x)}} \cdot df|_x = df|_{E_x} \cdot \pi|_{E_x}$, we obtain:

$$\frac{\partial G_1(u_1,u_2)}{\partial u_2} = df|_{E_x} \cdot \pi|_{E_x} \cdot \left(Id|_{F_x} + \frac{\partial G(v_1,v_2)}{\partial v_2} \right) \cdot \frac{\partial v_2}{\partial u_2} = df|_{E_x} \cdot \frac{\partial G(v_1,v_2)}{\partial v_2} \cdot \frac{\partial v_2}{\partial u_2}.$$

Applying (46) and the definition of dispersion, we deduce

$$\operatorname{disp}(G_1) \leq \|df|_{E_x}\| \cdot \operatorname{disp}(G) \cdot \|df^{-1}|_{F_{f(x)}}\|. \tag{48}$$

Thus, Inequality (47) follows, taking $c_1 := \max_{x \in M}\{\|df|_{E_x}\| \cdot \|df^{-1}|_{F_{f(x)}}\|\}$, ending the proof of Assertion 5.1. □

Assertion 5.2 *Let $\delta > 0$ be such that for all $x \in M$*

$$\exp_x : \{v \in T_xM : \|v\| \leq 3\delta\} \to B_{3\delta}(x) \subset M$$

is a diffeomorphism. For all $0 < \delta_0 < \delta$ there exists $0 < c' < 1/2$ with the following property: Assume that G is a Hadamard graph defined in $\mathbf{B}_\delta^{T_xM}(\mathbf{0})$ such that

$$\operatorname{disp} G < c'.$$

Assume that there exists $n \in \mathbb{N}$ and a graph G_n in $\mathbf{B}_\delta^{T_{f^n(x)}M}(\mathbf{0}) \subset T_{f^n(x)}M$ such that

$$\forall\ y = \exp_x(v_1 + v_2 + G(v_1,v_2)) \in B_{\delta_0}^n(x) \qquad \exists\ (u_1,u_2) \in \mathbf{B}_\delta^{E_{f^n(x)}}(\mathbf{0}) \times \mathbf{B}_\delta^{F_{f^n(x)}}(\mathbf{0})$$

where u_1 depends only on v_1 and

$$f^n(y) = \exp_{f^n(x)}(u_1 + u_2 + G_n(u_1,u_2)). \tag{49}$$

Then, the following inequality holds for all $y = \exp_x(v_1 + v_2 + G(v_1,v_2)) \in B_{\delta_0}^n(x)$:

$$\left\| \frac{\partial G_n}{\partial u_2}(u_1,u_2) \right\| \leq \left\| df^n|_{E_x} \right\| \cdot \operatorname{disp} G \cdot \left\| df^{-n}|_{F_{f^n(x)}} \right\|. \tag{50}$$

Proof: To simplify the notation, we do not write the exponential maps.

Equality (5.2) can be written as follows:

$$f^n(v_1 + v_2 + G(v_1,v_2)) = u_1 + u_2 + G_n(u_1,u_2), \tag{51}$$

where $(v_1, v_2) \in E_x \times F_x$, $(u_1, u_2) \in E_{f^n(x)} \times F_{f^n(x)}$, $G \in E_x$, $G_n \in E_{f^n(x)}$ and $y = v_1 + v_2 + G(v_1, v_2) \in B^n_{\delta_0}(x)$. Then:

$$u_2 = \pi_{F_{f^n(x)}} f^n(v_1 + v_2 + G(v_1, v_2)), \tag{52}$$

$$G_n(u_1, u_2) = -u_1 + \pi_{E_{f^n(x)}} f^n(v_1 + v_2 + G(v_1, v_2)). \tag{53}$$

Taking derivatives in Equality (52) with respect to v_2, with constant v_1, and noticing that $\pi|_{F_{f^n(x)}} \cdot df = df|_{F_x} \cdot \pi_{F_x}$, we obtain:

$$\frac{\partial u_2}{\partial v_2} = df^n|_{F_x} \pi_{F_x}(Id|_{F_x} + (\partial G/\partial v_2)) = df^n|_{F_x} = (df^{-n}|_{F_{f^n(x)}})^{-1}.$$

In the second equality above, we used that $G(v_1, v_2) \in E_x$ for all (v_1, v_2) (recall Definition 3.1 of Hadamard graphs). Since $df^n|_{F_x}$ is invertible, then the linear transformation $\partial u_2/\partial v_2$ is also invertible, and

$$\left(\frac{\partial u_2}{\partial v_2}\right)^{-1} = \frac{\partial v_2}{\partial u_2} = df^{-n}|_{F_{f^n(x)}}.$$

Now, we take derivatives in Equality (53) with respect to v_2 with constant v_1. We recall that, by hypothesis, u_1 depends only on v_1, but not on v_2. Besides, we notice that $\pi|_{E_{f^n(x)}} \cdot df^n = df^n|_{E_x} \cdot \pi|_{E^x}$. We obtain:

$$\frac{\partial G_n}{\partial u_2} \cdot \frac{\partial u_2}{\partial v_2} = df^n|_{E_x} \cdot \frac{\partial G}{\partial v_2}.$$

Thus:

$$\frac{\partial G_n}{\partial u_2} = df^n|_{E_x} \cdot \frac{\partial G}{\partial v_2} \cdot \frac{\partial v_2}{\partial u_2} = df^n|_{E_x} \cdot \frac{\partial G}{\partial v_2} \cdot df^{-n}|_{F_{f^n(x)}}.$$

So, after Definition 3.2 of $\mathrm{disp}(G)$ we deduce

$$\left\| \frac{\partial G_n}{\partial u_2} \right\| \leq \|df^n|_{E_x}\| \cdot \mathrm{disp}\, G \cdot \left\| df^{-n}|_{F_{f^n(x)}} \right\|,$$

proving Inequality (50). □

Assertion 5.3 .

There exists $n_0 \geq 1$ such that

$$-\sum_{i=1}^{k_n} p_i \log p_i \leq \frac{n\varepsilon}{5} + \frac{nH(\alpha^q, \mu_n)}{q} \quad \forall\ n \geq n_0,$$

where $\epsilon > 0$, α is a finite partition, $\alpha^n = \bigvee_{j=0}^n f^{-j}(\alpha)$, $q, n, k_n \in \mathbb{N}^+$, $k_n \leq \#(\alpha^n)$, $0 \leq p_i \leq 1$, $\sum_{i=1}^{k_n} p_i = 1$, $\mu_n := \frac{1}{n}\sum_{j=0}^{n-1}\sum_{i=1}^{k_n} p_i \delta_{f^j(x_i)}$ with $x_i \in X_i$ and $X_i \in \alpha^n$, and $H(\alpha^q, \mu_n) := -\sum_{A \in \alpha^q} \mu_n(A) \log \mu_n(A)$.

Proof: Denote $k := \#\alpha$. Construct the probability measure $\pi_n := \sum_{i=1}^{k_n} p_i \delta_{x_i}$. Then $\pi_n(X_i) = p_i \ \ \forall\ 1 \leq i \leq k_n$ and

$$H(\alpha^n, \pi_n) = -\sum_{i=1}^{k_n} p_i \log p_i$$

Fix $0 \leq l \leq q-1$. Since α^{n+l} is thinner than α^n, we have $H(\alpha^n, \pi_n) \leq H(\alpha^{n+l}, \pi_n)$. Thus

$$-\sum_{i=1}^{k_n} p_i \log p_i \leq H(\alpha^{n+l}, \pi_n), \tag{54}$$

where

$$\alpha^{n+l} = \vee_{j=0}^{n+l} f^{-j}\alpha = \Big(\vee_{j=0}^{l-1} f^{-j}\alpha \Big) \vee \Big(f^{-l}(\vee_{j=0}^{n} f^{-j}\alpha) \Big).$$

Besides, for any two partitions α and β, and for any probability measure ν we have $H(\alpha \vee \beta, \nu) \leq H(\alpha, \nu) + H(\beta, \nu)$. Therefore

$$H(\alpha^{n+l}, \pi_n) \leq \sum_{j=0}^{l-1} H(\alpha, f^{*j}\pi_n) + H(f^{-l}\alpha^n, \pi_n), \tag{55}$$

where the operator $f^* : \mathcal{P} \mapsto \mathcal{P}$ in the space of probability measures is defined by $f^*(\nu)(B) = \nu(f^{-1}(B))$ for any measurable set B.
Since $H(\alpha, \nu) \leq \log(\#(\alpha)) = \log k$ for any probability measure ν, and since $0 \leq l < q$, from Inequalities (54) and (55), we obtain:

$$-\sum_{i=1}^{k_n} p_i \log p_i \leq q \log k + H(\alpha^n, f^{*l}\pi_n).$$

If $n \geq (10\, q\, \log k) \,/\, \varepsilon$, then

$$-\sum_{i=1}^{k_n} p_i \log p_i \leq \frac{n\varepsilon}{10} + H(\alpha^n, f^{*l}\pi_n). \tag{56}$$

Now we write: $n = Nq + s,\ 0 \leq s \leq q-1$. We have

$$H(\alpha^n, f^{*l}\pi_n) \leq \sum_{h=0}^{N-1} H(\alpha^q, f^{*hq+l}\pi_n) + \sum_{j=Nq}^{Nq+s} H(\alpha, f^{*j+l}\pi_n)$$

Arguing as above:

$$\sum_{j=Nq}^{Nq+s} H(\alpha, f^{*j+l}\pi_n) \leq (s+1)\log k \leq q \log k \leq \frac{n\varepsilon}{10},$$

so Inequality (56) implies:

$$-\sum_{i=1}^{k_n} p_i \log p_i \leq \frac{n\varepsilon}{5} + \sum_{h=0}^{N-1} H(\alpha^q, f^{*hq+l}\pi_n)$$

Taking all the values of l such that $0 \leq l \leq q-1$ and adding the above bounds:

$$-q\sum_{i=1}^{k_n} p_i \log p_i \leq \frac{nq\varepsilon}{5} + \sum_{h=0}^{N-1}\sum_{l=0}^{q-1} H(\alpha^q, f^{*hq+l}\pi_n) \leq \frac{nq\varepsilon}{5} + \sum_{j=0}^{n-1} H(\alpha^q, f^{*j}\pi_n). \quad (57)$$

Recall that the entropy H of a partition with respect to a convex combination of probabilities, is larger than or equal to the convex combination of the entropies with respect to each of the probabilities. Since $\mu_n = \frac{1}{n}\sum_{j=1}^{n-1}\sum_{i=1}^{k_n} p_i \delta_{f^j(x_i)} = \frac{1}{n}\sum_{j=0}^{n-1} f^{*j}\pi_n$, we deduce $\frac{1}{n}\sum_{j=0}^{n-1} H(\alpha^q, f^{*j}\pi_n) \leq H(\alpha^q, \mu_n)$. Substituting in Inequality (57):

$$-q\sum_{i=1}^{k_n} p_i \log p_i \leq \frac{nq\varepsilon}{5} + nH(\alpha^q, \mu_n). \qquad \square$$

References

[CQ01] Campbell, J.; Quas, A.: *A generic C^1 expanding map has a singular S-R-B measure.* Commun. Math. Phys. **349** 2001 pp. 221-335.

[CE11] Catsigeras, E.; Enrich, H.: *SRB-like measures for C^0 dynamics.* Bull. Pol. Acad. Sci. Math. **59** 2011 pp. 151-164.

[K98] Keller, G.: *Equilibrium states in ergodic theory.* London Mathematics Society Student Texts.Cambridge Universtity Press 1998 and Corrections to Equilibrium states in ergodic theory in http://www.mi.uni-erlangen.de/∼keller/publications/equibook-corrections.pdf.

[LY85] Ledrappier, L.; Young, L.-S.: *The metric entropy of diffeomorphisms Part I: Characterization of measures satisfying Pesin's entropy formula.* Ann. of Math. **122** 1985 pp. 509-539.

[M81] Mañé, R.: *A proof of Pesin's formula.* Ergod. Th. & Dynam. Sys. 1981 **1**, 95-102 and *Errata to 'A proof of Pesin's formula'.* Ergod. Th. & Dynam. Sys. 1983 **3**, 159-160.

[P84] Pugh, C.: *The $C^{1+\alpha}$ hypothesis in Pesin theory.* Publ. Math. IHES, **59** 1984 pp. 143-161.

[PS89] Pugh, C., Shub, M.*Ergodic attractors* Trans. Amer. Math. Soc., **312** 1989 pp. 1-54.

[Q11] Qiu, H.: *Existence and uniqueness of SRB measure on C1 generic hyperbolic attractors.* Commun. Math. Phys. **302** 2011 pp. 345-357.

[R62] Rohlin, V.: *On the fundamental ideas of measure theory,* A. M. S. Transl. **1** 1962 pp. 1-52.

[R78] Ruelle, D.: *An inequality for the entropy of differentiable maps.* Bol. Soc. Bras. Mat. **9** 1978 pp. 83-87.

[ST12] Sun, W.; Tian, X.: *Dominated splitting and Pesin's entropy formula.* Disc. and Cont. Dyn. Syst. **32-4** 2012 pp. 1421-1434.